

ON THE DYNAMICAL SYMMETRIC ALGEBRA OF AGEING: LIE STRUCTURE, REPRESENTATIONS AND APPELL SYSTEMS

MALTE HENKEL^{*}, RENÉ SCHOTT[†], STOIMEN STOIMENOV[‡],
JÉRÉMIE UNTERBERGER[§]

The study of ageing phenomena leads to the investigation of a maximal parabolic subalgebra of $\mathfrak{conf}_3$ which we call $\mathfrak{alt}$. We investigate its Lie structure, prove some results concerning its representations and characterize the related Appell systems.

1. Introduction

Ageing phenomena occur widely in physics: glasses, granular systems or phase-ordering kinetics are just a few examples. While it is well-accepted that they display some sort of dynamical scaling, the question has been raised whether their non-equilibrium dynamics might possess larger symmetries than merely scale-invariance. At first sight, the noisy terms in the Langevin equations usually employed to model these systems might appear to exclude any non-trivial answer, but it was understood recently that provided the deterministic part of a Langevin equation is Galilei-invariant, then all observables can be exactly expressed in terms of multipoint correlation functions calculable from the deterministic part only.⁷ It is therefore of interest to study the dynamical symmetries of non-linear partial differential equations which extend dynamical scaling. In this context, the so-called Schrödinger algebra $\mathfrak{sch}$ has been shown to play an important rôle in phase-ordering kinetics. In what follows we shall restrict to one space dimension and we recall in figure 1 through a root diagram the definition of $\mathfrak{sch}$ as a parabolic subalgebra of the conformal algebra $\mathfrak{conf}_3$.⁶

^{*}lpm (cnrs umr 7556), université henri poincaré, bp 239, f - 54506 vandoeuvre-lès-nancy, france

[†]iecn and loria, université henri poincaré, bp 239, f - 54506 vandoeuvre-lès-nancy, france

[‡]lpm (cnrs umr 7556), université henri poincaré, bp 239, f - 54506 vandoeuvre-lès-nancy, france and institute of nuclear research and nuclear energy, bulgarian academy of sciences, 1784 sofia, bulgaria

[§]iecn, université henri poincaré, bp 239, f - 54506 vandoeuvre-lès-nancy, france




Figure 1. (a) Root diagram of the complexified conformal Lie algebra $(\mathfrak{conf}_3)_\mathbb{C}$ and the labelling of its generators. The double circle in the center denotes the Cartan subalgebra. The generators of the maximal parabolic subalgebras (b) $\mathfrak{sch}$ and (c) $\mathfrak{alt}$ are indicated by the filled dots.

2. A brief perspective on the algebra $\mathfrak{alt}$

There is a classification of semi-linear partial differential equations with a parabolic subalgebra of $\mathfrak{conf}_3$ as a symmetry.⁹ Here we shall study the abstract Lie algebra $\mathfrak{alt}$, which is the other maximal parabolic subalgebra of $\mathfrak{conf}_3$ (see figure 1) and its representations; we shall also see that, like the algebra $\mathfrak{sch}$, it can be embedded naturally in an infinite-dimensional Lie algebra $\mathcal{W}$ which is an extension of the algebra $\text{Vect}(S^1)$ of vector fields on the circle. Quite strikingly, we shall find on our way a 'no-go theorem' that proves the impossibility of a conventional extension of the embedding $\mathfrak{alt} \subset \mathfrak{conf}_3$.

2.1. The abstract Lie algebra $\mathfrak{alt}$

Elementary computations make it clear that (see figure 1 and $D = 2X_0 - N$)

$$\mathfrak{alt} = \langle V_+, D, Y_{-\frac{1}{2}} \rangle \ltimes \langle X_1, Y_{\frac{1}{2}}, M_0 \rangle := \mathfrak{g} \ltimes \mathfrak{h} \quad (1)$$

is a semi-direct product of $\mathfrak{g} \simeq \mathfrak{sl}(2, \mathbb{R})$ by a three-dimensional commutative Lie algebra $\mathfrak{h}$; the vector space $\mathfrak{h}$ is the irreducible spin-1 real representation of $\mathfrak{sl}(2, \mathbb{R})$, which can be identified with $\mathfrak{sl}(2, \mathbb{R})$ itself with the adjoint action. So one has the following

Proposition 2.1:

- (1) $\mathfrak{alt} \simeq \mathfrak{sl}(2, \mathbb{R}) \otimes \mathbb{R}[\varepsilon]/\varepsilon^2$, where ε is a 'Grassmann' variable;
- (2) $\mathfrak{alt} \simeq \mathfrak{p}_3$ where $\mathfrak{p}_3 \simeq \mathfrak{so}(2, 1) \ltimes \mathbb{R}^3$ is the relativistic Poincaré algebra in $(2+1)$ -dimensions.

Proof : The linear map $\Phi : \mathfrak{alt} \rightarrow \mathfrak{sl}(2, \mathbb{R}) \otimes \mathbb{R}[\varepsilon]/\varepsilon^2$ defined by

$$\Phi(V_+) = L_1, \quad \Phi(D) = L_0, \quad \Phi(Y_{-\frac{1}{2}}) = L_{-1}$$

$$\Phi(X_1) = \frac{1}{2}L_1^\varepsilon, \quad \Phi(Y_{\frac{1}{2}}) = L_0^\varepsilon, \quad \Phi(M_0) = L_{-1}^\varepsilon$$

is easily checked to be a Lie isomorphism. $\square$

In particular, the representations of $\mathfrak{alt} \simeq \mathfrak{p}_3$ are well-known since Wigner studied them in the 30'es.

2.2. Central extensions: an introduction

Consider any Lie algebra $\mathfrak{g}$ and an antisymmetric real two-form α on $\mathfrak{g}$. Suppose that its Lie bracket $[\cdot, \cdot]$ can be 'deformed' into a new Lie bracket $[\widetilde{\cdot}, \cdot]$ on $\widetilde{\mathfrak{g}} := \mathfrak{g} \times \mathbb{R}K$, where $[K, \mathfrak{g}] = 0$, by putting $[(X, 0), (Y, 0)] = ([X, Y], \alpha(X, Y))$. Then $\widetilde{\mathfrak{g}}$ is called a *central extension* of $\mathfrak{g}$. The Jacobi identity is equivalent with the nullity of the totally antisymmetric three-form $d\alpha : \Lambda^3(\mathfrak{g}) \rightarrow \mathbb{R}$ defined by

$$d\alpha(X, Y, Z) = \alpha([X, Y], Z) + \alpha([Y, Z], X) + \alpha([Z, X], Y).$$

Now we say that two central extensions $\mathfrak{g}_1, \mathfrak{g}_2$ of $\mathfrak{g}$ defined by α_1, α_2 are equivalent if $\mathfrak{g}_2$ can be gotten from $\mathfrak{g}_1$ by substituting $(X, c) \mapsto (X, c + \lambda(X))$ ($X \in \mathfrak{g}$) for a certain 1-form $\lambda \in \mathfrak{g}^*$, that is, by changing the non-intrinsic embedding of $\mathfrak{g}$ into $\widetilde{\mathfrak{g}}_1$. In other words, α_1 and α_2 are equivalent if $\alpha_2 - \alpha_1 = d\lambda$, where $d\lambda(X, Y) = \langle \lambda, [X, Y] \rangle$. The operator d can be made into the differential of a complex (called Chevalley-Eilenberg complex), and the preceding considerations make it clear that the classes of equivalence of central extensions of $\mathfrak{g}$ make up a vector space $H^2(\mathfrak{g}) = Z^2(\mathfrak{g})/B^2(\mathfrak{g})$, where Z^2 is the space of *cocycles* $\alpha \in \Lambda^2(\mathfrak{g}^*)$ verifying $d\alpha = 0$, and B^2 is the space of *coboundaries* $d\lambda$, $\lambda \in \mathfrak{g}^*$. We have the well-known

Proposition 2.2: *The Lie algebra $\mathfrak{alt}$ has no non-trivial central extension: $H^2(\mathfrak{alt}) = 0$.*

All this becomes very different when one embeds $\mathfrak{alt}$ into an infinite-dimensional Lie algebra.

2.3. Infinite-dimensional extension of $\mathfrak{alt}$

The Lie algebra $\text{Vect}(S^1)$ of vector fields on the circle has a long story in mathematical physics. It was discovered by Virasoro¹⁰ in the 70'es that

$\text{Vect}(S^1)$ has a one-parameter family of central extensions which yield the so-called *Virasoro algebra*

$$\mathfrak{vir} := \text{Vect}(S^1) \oplus \mathbb{R}K = \langle (L_n)_{n \in \mathbb{Z}}, K \rangle \quad (2)$$

with Lie brackets

$$[K, L_n] = 0, \quad [L_n, L_m] = (n - m)L_{n+m} + \delta_{n+m,0} c n(n^2 - 1)K \quad (c \in \mathbb{R})$$

When $c = 0$, one retrieves $\text{Vect}(S^1)$ by identifying the (L_n) with the usual Fourier basis $(e^{in\theta} d\theta)_{n \in \mathbb{Z}}$ of periodic vector fields on $[0, 2\pi]$, or with $-z^{n+1} \frac{d}{dz}$ with $z := e^{i\theta}$. Note in particular that $\langle L_{-1}, L_0, L_1 \rangle$ is isomorphic to $\mathfrak{sl}(2, \mathbb{R})$, and that the Virasoro cocycle restricted to $\mathfrak{sl}(2, \mathbb{R})$ is 0, as should be (since $\mathfrak{sl}(2, \mathbb{R})$ has no non-trivial central extensions).

It is tempting to embed $\mathfrak{alt} \simeq \mathfrak{sl}(2, \mathbb{R}) \otimes \mathbb{R}[\varepsilon]/\varepsilon^2$ into the Lie algebra

$$\mathcal{W} := \text{Vect}(S^1) \otimes \mathbb{R}[\varepsilon]/\varepsilon^2 = \langle L_n \rangle_{n \in \mathbb{Z}} \ltimes \langle L_n^\varepsilon \rangle_{n \in \mathbb{Z}}, \quad (3)$$

with Lie brackets

$$[L_n, L_m] = (n - m)L_{n+m}, \quad [L_n, L_m^\varepsilon] = (n - m)L_{n+m}^\varepsilon, \quad [L_n^\varepsilon, L_m^\varepsilon] = 0.$$

These brackets come out naturally putting $\mathcal{W}$ in the 2×2 -matrix form

$$L_n \mapsto \begin{pmatrix} L_n & 0 \\ 0 & L_n \end{pmatrix}, \quad L_n^\varepsilon \mapsto \begin{pmatrix} 0 & L_n \\ 0 & 0 \end{pmatrix} \quad (4)$$

leading to straightforward generalizations (there exists a deformation of $\mathfrak{so}$ that can be represented as upper-triangular 3×3 Virasoro matrices⁸).

In terms of the standard representations of $\text{Vect}(S^1)$ as modules of α -densities $\mathcal{F}_\alpha = \{u(z)(dz)^\alpha\}$ with the action

$$f(z) \frac{d}{dz} (u(z)(dz)^\alpha) = (fu' + \alpha f'u)(z)(dz)^\alpha, \quad (5)$$

we have

Proposition 2.3: $\mathcal{W} \simeq \text{Vect}(S^1) \ltimes \mathcal{F}_{-1}$.

There are two linearly independent central extensions of $\mathcal{W}$:

1. the natural extension to $\mathcal{W}$ of the Virasoro cocycle on $\text{Vect}(S^1)$, namely $[\cdot, \cdot] = \widetilde{[\cdot, \cdot]}$ except for $[\widetilde{L_n}, \widetilde{L_{-n}}] = n(n^2 - 1)K + 2nL_0$. In other words, $\text{Vect}(S^1)$ is centrally extended, but its action on $\mathcal{F}_{-1}$ remains unchanged;

2. the cocycle ω which is zero on $\Lambda^2(\text{Vect}(S^1))$ and $\Lambda^2(\mathcal{F}_{-1})$, and defined by $\omega(L_n, L_m^\varepsilon) = \delta_{n+m,0} n(n^2 - 1)K$ on $\text{Vect}(S^1) \times \mathcal{F}_{-1}$.

A natural related question is: can one deform the extension of $\text{Vect}(S^1)$ by the $\text{Vect}(S^1)$ -module $\mathcal{F}_{-1}$? The answer is: no, thanks to the triviality³⁵

of the cohomology space $H^2(\text{Vect}(S^1), \mathcal{F}_{-1})$. Hence, any Lie algebra structure $\widetilde{[\cdot, \cdot]}$ on the vector space $\text{Vect}(S^1) \oplus \mathcal{F}_{-1}$ such that

$$[(X, \phi), (Y, \psi)] = ([X, Y]_{\text{Vect}(S^1)}, \text{ad}_{\text{Vect}(S^1)} X \cdot \psi - \text{ad}_{\text{Vect}(S^1)} Y \cdot \phi + B(X, Y))$$

is isomorphic to the Lie structure of $\mathcal{W}$ (where B is an antisymmetric two-form on $\text{Vect}(S^1)$). So one may say that $\mathcal{W}$ and its central extensions are natural objects to look at.

2.4. Some results on representations of $\mathcal{W}$

We now state two results which may deserve deeper thoughts and will be developed in the future.

Proposition 2.4 ('no-go theorem'): *There is no way to extend the usual representation of $\mathfrak{alt}$ as conformal vector fields into an embedding of $\mathcal{W}$ into the Lie algebra of vector fields on $\mathbb{R}^3$.*

Proposition 2.5: *The infinite-dimensional extension $\mathcal{W}$ of the algebra $\mathfrak{alt}_1$ is a contraction of a pair of commuting Virasoro algebras $\mathfrak{vir} \oplus \overline{\mathfrak{vir}} \rightarrow \mathcal{W}$. In particular, we have the explicit differential operator representation*

$$\begin{aligned} L_n &= -t^{n+1} \partial_t + (n+1)t^n r \partial_r - (n+1)xt^n - n(n+1)\gamma t^{n-1}r \\ L_n^\varepsilon &= -t^{n+1} \partial_r - (n+1)\gamma t^n \end{aligned} \quad (6)$$

where x and γ are parameters and $n \in \mathbb{Z}$.

3. Appell systems

Definition 3.1. *Appell polynomials $\{h_n(x); n \in \mathbb{N}\}$ on $\mathbb{R}$ are usually characterized by the two conditions*

- $h_n(x)$ are polynomials of degree n ,
- $Dh_n(x) = nh_{n-1}(x)$, where D is the usual derivation operator.

Interesting examples are furnished by the shifted moment sequences

$$h_n(x) = \int_{-\infty}^{\infty} (x+y)^n p(dy)$$

where p is a probability measure on $\mathbb{R}$ with all moments finite.

This definition generalizes to higher dimensions. On non-commutative algebraic structures, the shifting corresponds to left or right multiplication and in general, $\{h_n\}$ is not a family of polynomials. We shall call it *Appell systems*.²

Appell systems of the Schrödinger algebra $\mathfrak{sch}$ have been investigated¹ but the algebra $\mathfrak{alt}$ requires a specific study. $\mathfrak{alt}$ has the following Cartan decomposition:

$$\mathfrak{alt} = \mathfrak{P} \oplus \mathfrak{K} \oplus \mathfrak{L} = \{Y_1, X_1\} \oplus \{Y_0, X_0\} \oplus \{Y_{-1}, X_{-1}\} \quad (7)$$

and there is a one to one correspondence between the subalgebras $\mathfrak{P}$ and $\mathfrak{L}$. Write $X \in \mathfrak{alt}$ in the form $X = \sum_{i=1}^6 \alpha_i b_i$, where $\{b_i, i = 1, \dots, 6\}$ is a basis of $\mathfrak{alt}$. The α_i are called coordinates of the first kind.

Here we use the basis $b_1 = Y_1$, $b_2 = X_1$, $b_3 = Y_0$, $b_4 = X_0$, $b_5 = Y_{-1}$, $b_6 = X_{-1}$. Let ALT be the simply connected Lie group corresponding to $\mathfrak{alt}$. Group elements in a neighbourhood of the identity can be expressed as

$$e^X = e^{A_1 b_1} \dots e^{A_6 b_6}$$

The A_i are called *coordinates of the second kind*.

Referring to the decomposition (7), we specialize variables, writing V_1, V_2, B_1, B_2 for A_1, A_2, A_5, A_6 respectively. Basic for our approach is to establish the partial group law: $e^{B_1 Y_{-1} + B_2 X_{-1}} e^{V_1 Y_1 + V_2 X_1} = ?$. We get

$$B_1 Y_{-1} + B_2 X_{-1} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ -B_2 & 0 & -B_1 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -B_2 & 0 \end{pmatrix}, \quad V_1 Y_1 + V_2 X_1 = \begin{pmatrix} 0 & V_2 & 0 & V_1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & V_2 \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (8)$$

and finally:

$$e^{B_1 Y_{-1} + B_2 X_{-1}} e^{V_1 Y_1 + V_2 X_1} = \begin{pmatrix} 1 & V_2 & 0 & V_1 \\ -B_2 & 1 - B_2 V_2 & -B_1 & -B_2 V_1 - B_1 V_2 \\ 0 & 0 & 1 & V_2 \\ 0 & 0 & -B_2 & 1 - B_2 V_2 \end{pmatrix} \quad (9)$$

Proposition 3.1. *In coordinates of the second kind, we have the Leibnitz formula,*

$$\begin{aligned} g(0, 0, 0, 0, B_1, B_2) g(V_1, V_2, 0, 0, 0, 0) &= g(A_1, A_2, A_3, A_4, A_5, A_6) = \\ g\left(\frac{B_1 V_2^2 + V_1}{(1 - B_2 V_2)}, \frac{V_2}{(1 - B_2 V_2)}, -2 \frac{B_1 V_2 + B_2 V_1}{(1 - B_2 V_2)}, 2 \ln(1 - B_2 V_2), \right. & \\ \left. \frac{B_1 - 2 B_1 B_2 V_2 - B_2^2 V_1}{(1 - B_2 V_2)^2}, \frac{B_2}{(1 - B_2 V_2)}\right) & \end{aligned} \quad (10)$$

Now we are ready to construct the representation space and basis-the canonical Appell system. To start, define a vacuum state Ω . The elements Y_1, X_1 of $\mathfrak{P}$ can be used to form basis elements

$$|jk\rangle = Y_1^j X_1^k \Omega, j, k \geq 0 \quad (11)$$

of a Fock space $\mathfrak{F} = \text{span}\{|jk\rangle\}$ on which Y_1, X_1 act as raising operators, Y_{-1}, X_{-1} as lowering operator and Y_0, X_0 as multiplication with the constants γ, x (up to the sign) correspondingly. That is,

$$\begin{aligned} Y_1\Omega &= |10\rangle, X_1\Omega = |01\rangle \\ Y_{-1}\Omega &= 0, X_{-1}\Omega = 0 \\ Y_0\Omega &= -\gamma|00\rangle, X_0\Omega = -x|00\rangle \end{aligned} \quad (12)$$

The goal is to find an abelian subalgebra spanned by some selfadjoint operators acting on representation space, just constructed. Such a two-dimensional subalgebra can be obtained by an appropriate “turn” of the plane $\mathfrak{P}$ in the Lie algebra, namely via the adjoint action of the group element formed by exponentiating X_{-1} . The resulting plane, $\mathfrak{P}_\beta$ say, is abelian and is spanned by

$$\begin{aligned} \bar{Y}_1 &= e^{\beta X_{-1}} Y_1 e^{-\beta X_{-1}} = Y_1 - 2\beta Y_0 + \beta^2 Y_{-1} \\ \bar{X}_1 &= e^{\beta X_{-1}} X_1 e^{-\beta X_{-1}} = X_1 - 2\beta X_0 + \beta^2 X_{-1} \end{aligned} \quad (13)$$

Next we determine our canonical Appell systems. We apply the Leibniz formula (10) with $B_1 = 0, B_2 = \beta, V_1 = z_1, V_2 = z_2$ and (12). This yields

$$\begin{aligned} e^{z_1 \bar{Y}_1} e^{z_2 \bar{X}_1} \Omega &= e^{\beta X_{-1}} e^{z_1 Y_1} e^{z_2 X_1} e^{-\beta X_{-1}} \Omega = e^{\beta X_{-1}} e^{z_1 Y_1} e^{z_2 X_1} \Omega = \\ &= e^{\frac{z_1 Y_1}{(1-\beta z_2)^2}} e^{\frac{z_2 X_1}{(1-\beta z_2)}} e^{\frac{2\gamma\beta z_1}{(1-\beta z_2)}} (1-\beta z_2)^{-2x} \Omega \end{aligned} \quad (14)$$

To get the generating function for the basis $|jk\rangle$ set in equation (14)

$$v_1 = \frac{z_1}{(1-\beta z_2)^2}, v_2 = \frac{z_2}{(1-\beta z_2)} \quad (15)$$

Substituting throughout, we have

Proposition 3.2. *The generating function for the canonical Appell system, $|jk\rangle = Y_1^j X_1^k \Omega$ is*

$$\begin{aligned} e^{v_1 Y_1 + v_2 X_1} \Omega &= \exp(y_1 \frac{v_1}{(1+\beta v_2)^2}) \exp(y_2 \frac{v_2}{(1+\beta v_2)}) \exp(-\frac{2\gamma\beta v_1}{(1-\beta v_2)}) \\ &\quad (1+\beta v_2)^{-2x} \Omega \end{aligned} \quad (16)$$

where we identify $\bar{Y}_1 \Omega = y_1 \cdot 1$ and $\bar{X}_1 \Omega = y_2 \cdot 1$ in the realization as function of y_1, y_2 .

With $v_1 = 0$, we recognize the generating function for the Laguerre polynomials, while $v_2 = 0$ reduces to the generating function for Hermite polynomials.

References

1. P. Feinsilver, Y. Kocik and R. Schott, *Representations of the Schrödinger Algebra and Appell Systems*, Progress of Physics, **52** (2004) 343-359.
2. P. Feinsilver and R. Schott, *Algebraic Structures and Operator Calculus, Vol.3: Representations of Lie Groups*. Kluwer Academic Publishers, Dordrecht, 1993.
3. D. B. Fuks, *Cohohomology of infinite-dimensional Lie algebras*, Contemporary Soviet Mathematics, Consultants Bureau, New York, 1986.
4. M. Henkel, R. Schott, S. Stoimenov and J. Unterberger, *On the dynamical symmetric algebra of ageing: Lie structure, representations and Appell systems*. Prépublication Institut Elie Cartan, 2005.
5. L. Guieu, C. Roger, preprint, available on http://www.math.univ-montp2.fr/~guieu/The_Virasoro_Project/Phase1/.
6. M. Henkel, J. Unterberger, *Schrödinger-invariance and space-time symmetries*, Nuclear Physics **B660** (2003) 407-435.
7. A. Picone and M. Henkel, *Local scale-invariance and ageing in noisy systems*, Nuclear Physics **B688** (2004) 217-265.
8. C. Roger and J. Unterberger, in progress (2005).
9. S. Stoimenov and M. Henkel, *Dynamical symmetries of semi-linear Schrödinger and diffusion equations*, Nuclear Physics **B723** (2005) 205-233.
10. M. Virasoro, Phys. Rev. **D1**, 2933-2936 (1970).